

Asymptotics of Plancherel measures for $GL(n, q)$.

A. Dudko

Abstract

We introduce the Plancherel measures on the sets of partition collections $\{\Lambda_\phi\}$, which parameterize irreducible representations of $GL(n, q)$. Consider the Plancherel random collection $\{\Lambda_\phi\}$. We prove that as $n \rightarrow \infty$, the random partitions Λ_ϕ converge in finite dimensional distribution to independent random variables. We give explicit formulas for the corresponding limit distributions.

1. Introduction. Let $GL(n, q)$ be the group of all invertible $n \times n$ matrices over $\mathbb{F}_q$, where q is a prime power. Denote by Φ the set of all irreducible polynomials over $\mathbb{F}_q$ with unit highest coefficient and nonzero constant term. The irreducible representations of $GL(n, q)$ are parameterized by collections $\Lambda = \{\Lambda_\phi\}$ of partitions, where $\phi \in \Phi$ and

$$|\Lambda| = \sum_{\phi} |\Lambda_\phi| \deg \phi = n.$$

Denote by $\mathbb{L}_n$ the set of collections $\Lambda = \{\Lambda_\phi\}$ with $|\Lambda| = n$. Define the Plancherel measure μ_n on $\mathbb{L}_n$ as follows:

$$\mu_n(\Lambda) = \frac{d_\Lambda^2}{|GL(n, q)|}, \quad (1)$$

where d_Λ is the degree of the irreducible representation of $GL(n, q)$ corresponding to $\Lambda \in \mathbb{L}_n$. The Plancherel measure defines relative dimensions of regular representation isotopic components.

For $0 < v < q$ and $q > 1$ denote the measure $M_{v,q}$ on the set of all partitions of all numbers as follows:

$$M_{v,q}(\lambda) = \prod_{r=1}^{\infty} (1 - vq^{-r})^r q^{2n(\lambda') + |\lambda|} \prod_{x \in \lambda} (q^{h(x)} - 1)^{-2} v^{|\lambda|}, \quad (2)$$

where $h(x)$ is the hook-length and $n(\lambda') = \sum_i \lambda_i(\lambda_i - 1)/2$. We show, that $M_{v,q}$ is probabilistic. As we shall see, the measure $M_{v,q}$ is a specialization of the Schur measure (see [7]).

Let Λ be a random value in $\mathbb{L}_n$ distributed by (1). Then each $\phi \in \Phi$ denotes the random value Λ_ϕ in the set of partitions. We consider asymptotical behavior of Λ_ϕ with increase of n . The main result of this paper is the following

Theorem 1. *Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be arbitrary partitions. Let $\phi_1, \phi_2, \dots, \phi_k \in \Phi$ be different polynomials. Then there exists the limit:*

$$\lim_{n \rightarrow \infty} \mu_n(\{\Lambda \in \mathbb{L}_n : \Lambda_{\phi_j} = \lambda_j, j = 1, \dots, k\}) = \prod_{j=1}^k M_{1, q^{\deg \phi_j}}(\lambda_j). \quad (3)$$

The proof is based on the cycle index technic (see [3], [4]). Asymptotical behavior of Plancherel measures for symmetric groups is well studied. See for example [2].

I am grateful to N.I. Nessonov for the statement of the problem and constant attention to my work. I also would like to thank A. Borodin for the important remarks on the result of this paper. After this paper was posted to arxiv, the author knew from J. Fulman, that the results of this paper with proofs are contained in [5].

2. Notions. Make some preliminaries. Let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ be arbitrary partition. By definition, $n(\lambda) = \sum \lambda'_j(\lambda'_j - 1)/2$. The hook-length in $x = (i, j) \in \lambda$ is equal to

$$h(x) = \lambda_i + \lambda'_j - i - j + 1. \quad (4)$$

Since $n(\lambda) = \sum_{(i,j) \in \lambda} (\lambda'_j - i)$, then

$$\sum_{x \in \lambda} h(x) = n(\lambda) + n(\lambda') + |\lambda|. \quad (5)$$

From [6], I.3, ex. 2, using (5), one can deduce the following specialization formula for the Schur functions:

$$s_\lambda(q^{-1}, q^{-2}, \dots) = q^{n(\lambda')} \prod_{x \in \lambda} (q^{h(x)} - 1)^{-1} \quad (\text{see also [6], IV.6}). \quad (6)$$

By definition of the Schur measure $\mathfrak{M}$,

$$\mathfrak{M}(\lambda) = \prod_{i,j} (1 - x_i y_j) s_\lambda(x) s_\lambda(y). \quad (7)$$

From (2), (6) and (7) the next statement follows.

Proposition 2. *Let $0 < v < q$ and $q > 1$. If we set $x_i = q^{-i}$, $y_j = vq^{1-j}$ then the Schur measure $\mathfrak{M}$ specializes to $M_{v,q}$.*

By the Cauchy identity for the Schur functions, it follows that $M_{v,q}$ is probabilistic. Recall now the formula for the degree of the irreducible representation of $GL(n, \mathbb{F}_q)$ corresponding to $\Lambda = \{\Lambda_\phi\}$:

$$d_\Lambda = \prod_{i=1}^n (q^i - 1) \prod_{\phi} q^{n(\lambda') \deg \phi} \prod_{x \in \lambda} (q^{h(x) \deg \phi} - 1)^{-1} \quad (\text{see [6], IV.6}). \quad (8)$$

By (1), (6) and (8), one has:

$$\mu_n(\Lambda) = q^{-n(n-1)/2} \prod_{i=1}^n (q^i - 1) \prod_{\phi} s_{\Lambda_\phi}^2(q^{-\deg \phi}, q^{-2\deg \phi}, \dots). \quad (9)$$

3. The proof of the main result. Let $0 < v < 1$ and $q > 1$. Introduce the measure $P_{v,q}$ on $\mathbb{L} = \bigcup_{n=0}^{\infty} \mathbb{L}_n$ as follows:

$$P_{v,q}(\Lambda) = \prod_{r=0}^{\infty} (1 - vq^{-r}) q^{n(n+1)/2} \prod_{i=1}^n (q^i - 1)^{-1} v^n \mu_n(\Lambda) \quad \text{for } \Lambda \in \mathbb{L}_n. \quad (10)$$

It follows from the Euler's identity (see [1], page 19), that

$$\sum_{n=0}^{\infty} q^{n(n+1)/2} \prod_{i=1}^n (q^i - 1)^{-1} v^n = \prod_{r=0}^{\infty} (1 - vq^{-r})^{-1}. \quad (11)$$

Therefore, $P_{v,q}$ is probabilistic. For each $A \subset \mathbb{L}$ one has:

$$P_{v,q}(A) \prod_{r=0}^{\infty} (1 - vq^{-r})^{-1} = \sum_{n=0}^{\infty} q^{n(n+1)/2} \prod_{i=1}^n (q^i - 1)^{-1} v^n \mu_n(A \cap \mathbb{L}_n). \quad (12)$$

The right side of (12) is the Taylor expansion for the left side in $|v| < 1$. Consequently, for each n ,

$$\prod_{i=1}^n (1 - q^{-i})^{-1} \mu_n(A \cap \mathbb{L}_n) = [v^n] \prod_{r=0}^{\infty} (1 - vq^{-r})^{-1} P_{v,q}(A), \quad (13)$$

where $[v^n]f(v)$ stands for the coefficient of the n -th power in the Taylor expansion for f . Further, by (1), (2), (8) and (10), for each $\Lambda \in \mathbb{L}_n$,

$$\prod_{r=0}^{\infty} (1 - vq^{-r})^{-1} P_{v,q}(\Lambda) = \prod_{\phi} \left(M_{v^{deg \phi}, q^{deg \phi}}(\Lambda_{\phi}) \prod_{r=1}^{\infty} (1 - (vq^{-r})^{deg \phi})^{-r} \right). \quad (14)$$

In particular, as $P_{v,q}$ and $M_{v,q}$ are probabilistic, one has:

$$\prod_{r=0}^{\infty} (1 - vq^{-r})^{-1} = \prod_{\phi} \prod_{r=1}^{\infty} (1 - (vq^{-r})^{deg \phi})^{-r}. \quad (15)$$

Now fix $k \in \mathbb{N}$, arbitrary partitions $\lambda_j \in \mathbb{Y}$ and distinct polynomials $\phi_j \in \Phi, j = 1, \dots, k$. Using (13-15) we get:

$$\begin{aligned} \prod_{i=1}^n (1 - q^{-i})^{-1} \mu_n(\{\Lambda : \Lambda_{\phi_j} = \lambda_j, j = 1, \dots, k\} \cap \mathbb{L}_n) = \\ [v^n] \prod_{r=0}^{\infty} (1 - vq^{-r})^{-1} \prod_{j=1}^k M_{v^{deg \phi_j}, q^{deg \phi_j}}(\lambda_j). \end{aligned} \quad (16)$$

To finish the proof of the theorem 1 we need the following statement (see [4], paragraph 2.2, lemma 2):

Lemma 3. *Let $f(v)$ be a holomorphic in $|v| < 1$ function, such that the Taylor series of f around 0 converges at $v = 1$. Then the next limit exists:*

$$\lim_{n \rightarrow \infty} [v^n]f(v)(1 - v)^{-1} = f(1).$$

The proof follows from the identity $(1 - v)^{-1} = 1 + v + v^2 + \dots$ for $|v| < 1$. Using the lemma (3) for

$$f(v) = \prod_{r=1}^{\infty} (1 - vq^{-r})^{-1} \prod_{j=1}^k M_{v^{deg \phi_j}, q^{deg \phi_j}}(\lambda_j)$$

one obtains from (16):

$$\lim_{n \rightarrow \infty} \mu_n(\{\Lambda : \Lambda_{\phi_j} = \lambda_j, j = 1, \dots, k\} \cap \mathbb{L}_n) = \prod_{j=1}^k M_{1, q^{deg \phi_j}}(\lambda_j).$$

References

- [1] Andrews, The theory of partitions. Encyclopedia of Mathematics and its applications, Vol. 2. Addison-Wesley Publishing Co., Reading, Mass.-London-Amsterdam, 1976.
- [2] Borodin A., Okounkov A., Olshanski G., *Asymptotics of Plancherel measures for symmetric groups*, J. Amer. Math. Soc. 13 (2000), no. 3, 481–515, math.CO/9905032.
- [3] Fulman J., *Probability in the classical groups over finite fields: symmetric functions, stochastic algorithms and cycle indeces*, Ph.D. Thesis, Harvard University, 1997.
- [4] Fulman J., *Random matrix theory over finite fields*, arXiv:math/0003195.
- [5] Fulman J., *Convergence rates of random walk on irreducible representations of finite groups*, J. Theoret. Probab. Vol. 21, March 2008, arXiv:math/0607399.
- [6] Macdonald I., Symmetric functions and Hall Polynomials. Clarendon Press, Oxford. 1995.
- [7] Okounkov A., *Infinite wedge and random partitions*, Selecta Math., New Ser., 7 (2001), 57–81, math.RT/9907127.

Mathematics Division, Institute for Low Temperature Physics and Engineering, 47 Lenin ave., Kharkov 61103, Ukraine
e-mail: artemdudko@rambler.ru